Supplementary Table 2. Demographics and laboratory findings of the enrolled patients on admission

Variable	Normal diastolic function (n=83)	Diastolic dysfunction (n=121)	P-value
Age (yr)	66.0 (54.5–75.5)	74.0 (63.0–79.0)	0.004
Male sex	51 (61.4)	56 (46.3)	0.045
Hypertension	45 (54.2)	86 (71.1)	0.017
Diabetes	22 (26.5)	53 (43.8)	0.013
Heart failure	2 (2.4)	10 (8.3)	0.128
Atrial fibrillation	7 (8.4)	27 (22.3)	0.009
Coronary artery occlusive disease	7 (8.4)	26 (21.5)	0.013
Cerebrovascular accident	17 (20.5)	29 (24.0)	0.558
Chronic kidney disease	16 (19.3)	22 (18.2)	0.844
Malignancy	21 (25.3)	30 (24.8)	0.934
Sepsis origin			0.183
Respiratory	17 (20.5)	20 (16.5)	
Urinary tract	28 (33.7)	53 (43.8)	
Intra-abdominal infection	18 (21.7)	25 (20.7)	
Soft tissue infection	8 (9.6)	16 (13.2)	
Other/unknown	12 (14.5)	7 (5.8)	
Antibiotic administration time (min)	55.0 (35.0–75.5)	57.0 (39.0–74.0)	0.428
Age-adjusted Charlson Comorbidity Index	4.0 (1.5–5.0)	5.0 (3.0–6.0)	0.002
SOFA score	9.0 (7.0–11.0)	9.0 (7.0–12.0)	0.312
Outcome			
Hospital day	15.0 (9.0–25.5)	16.0 (9.0–30.0)	0.776
30-Day mortality	6 (7.2)	19 (15.7)	0.084
Vasopressor-free day	12.0 (5.0–20.5)	13.0 (6.0–25.0)	>0.999
ICU admission	48 (57.8)	71 (58.7)	>0.999
ICU day	3.0 (2.0–7.0)	5.0 (2.0–8.5)	0.314
Mechanical ventilation	16 (19.3)	35 (28.9)	0.162
Mechanical ventilation day	7.0 (2.0–13.5)	7.0 (3.5–15.5)	0.986

Values are presented as median (interquartile range), or number (%).
SOFA: Sequential Organ Failure Assessment; ICU: intensive care unit.